Factors associated with oro-facial injuries among children in Al-baha, Saudi Arabia

¹TO Lawoyin, ²DO Lawoyin, and ³JO Lawoyin

¹Department of Community Medicine, ²Department of Oral Pathology, College of Medicine, Univsersity College Hospital, Ibadan, Nigeria. ³Department of Oral Surgery, King Fahd Hospital atAl-baha, Kingdom of Saudi Arabia

Summary

This study was prompted by the dearth of injury data among children in Saudi Arabia. Data were collected prospectively on all children 0-15 years who presented to the King Fahad Hospital at Al Baha with oro-facial injuries over a period of five consecutive years. These records were analyzed for age, gender. seasonal distribution, cause, location and types of injuries. The mean age of the injured children was 6.2 ± 3.7 years. Falls, motor vehicle accidents followed by direct blunt trauma were responsible for over 97 percent of all oro-facial injuries in children, and over 60 percent of all injuries were due to falls. The forehead, eye, periorbital areas and the mouth accounted for about two- thirds of all injury sites. Most injuries 356 (75.3%) occurred during the summer months while only 117 (24.7%) occurred during the winter months. The difference was significant (P<0.0001). The large majority of injuries, which occurred in children under the age of 6 years, took place at home, while road traffic accidents were the commonest cause of injury in the older children 6 years and above. Following logistic regression analysis, age (6 years and under) was associated with increased risk of injury while the winter months were associated with a lower risk. In addition, the study found no statistically significant difference in the prevalence by sex. Having a safe home environment, health education programmes directed towards the caretakers of young children and legislation in support of seat belts will go a long way towards reducing injuries among children in this community. A multi-disciplinary clinical management approach is envisaged due to the different injury sites and facilities would need to be equipped to take care of all types of orofacial injuries.

Keywords: Oro facial injuries, children, associated factors, Saudi Arabia.

Résumé

Cette etude a ete motivee par le manqué de donnees des blessures parmi les enfants-en Arabic Saoudite. Les donnees out ete collectees prospectivement sur tous les enfants de o a 15 ans qui se sont presentes a l'hospital king Fahad a Alo Baha avec des blessures orofaciales sur une periode de cinq annees consecutives. Ces rapports ont ete analyses suivant l'age, le sese, la raison de distribution, cause localite et types de blessures. L'age moyen des enfants blesses etait de 6,2,+3,7 ans. Les tombees, accidents de voitures suivi par des traumas direct et brusques etaient responsible de plus de 97% et brusques etaient responsible de plus de 97% et plus de 60% des blessures etaient causes par les chutes. Le front, les yeux, les zones peri-orbitales et la bouche convrent pres du deux tiers de tous les sites de blessures. La plupart des blessures 356 (75,3% ont eu lieu en ete alors que seulement 117 (24,7%) en hiver. La difference etait significative(P<0,0001). La grande majorite des blessures chez les

Correspondence: Dr Taiwo Lawoyin, Department of Community Medicine, University College Hospital, Ibadan, Nigeria.

enfants de moins de 6 ans a en liena la maison, alors les accidents de route etaient la cause commune de blessures chez les enfants de plus de 6 ons. L'analyse dem regression multiple, l'age (moins de 6 ona) etait associe a un risqué intense de blessure alors que les mois d'hiver etaient associes a un risqué peu eleve.. En plus, l'etude ne montre aucune difference significative dans la prevalence du sex. Avec un environnement sans danger les programmes de sante d'education tournes vers les gardiens des jeunes enfants et une legislation en support des ceintures de securite auront une longue portee sur la reductionm des blessures d'enfants dans cette communaute. Une gestion clinique multidisciplinaire est envisagee a cause des differents sites de blessures, et le besoin d'equiper (d'ameliorer) les facilites pour prendre soins de tous les cas de blessures oro – facials.

Introduction

Injuries are the commonest cause of death in young children in the USA and studies has found maxillo-facial injuries to be relatively common in children because of their greater cranialmass-to-body ratio [1]. Etiologies of such injuries are diverse among which include abuse and neglect, sports, motor vehicle accidents, firearms, attack by animals, burns and interpersonal violence [2-9]. Sex differences have been noted, with some studies describing significantly more injuries in males, while others found no significant sex difference [1,10-13]. Data on trauma to the orofacial region are important because of the untoward sequel and effect on development. Dentofacial injuries, which occur for example, before the primary dentition erupts, can result in damages to the primary and permanent dentition [14]. The effect of orofacial injuries can be quite serious, even life threatening if prompt and adequate care is not received [12,14-16]. Knowledge of the types and pattern of injury and factors associated with these injuries are therefore useful when preparing and implementing preventive strategies. They are also necessary for health care planning purposes at facility level.

Oro facial injuries in children are common in Saudi Arabia but there is virtually no literature on such injuries especially from the Al Baha region. Too few reports are available also on etiology and pattern of orofacial injuries in children from Saudi Arabia. This prompted the study, which looks at etiology and pattern of, as well as some factors associated with orofacial injury among children in AL-Baha region of Saudi Arabia.

Materials and methods

Al-Baha region is situated in the southwestern part of Saudi Arabia and has a population of about 500,000. The King Fahad Hospital, a 450-bedded hospital receives referrals from seven smaller hospitals as well as from 24 primary health care clinics within its catchments area. The vast majority of orofacial injuries and other accidents are referred to this hospital. The data collected is therefore a fair representation of what occurs at community level. Data were collected prospectively on all children who visited the King Fahad Hospital emergency room and the oral surgery clinic records from April 1990 to March 1995. These records were analyzed for age, gender, cause, location, seasonal distribution and types of injuries. All children 0-15 years who presented to the emergency room or clinic with orofacial injuries were included in the study. Data were entered in the computer and analyzed using EPI INFO version 6.0[17]. It was exported to Systat program for multiple regression analysis [18]. Permission to carry out the study was obtained from the Chief of Surgery, King Fahd Hospital, Al Baha.

Results

Four hundred and seventy-three children were seen and treated for orofacial injuries in the emergency room and oral surgery clinic over a 5-year period. Only one hundred and thirty (27.5 percent) had injuries which, necessitated admission. Admission days ranged from 1-33 days with a mean of 5.6 days. Injuries, which were due to road traffic accidents, had the longest mean admission days (7.5 days).

Age distribution

Mean age for all children was 6.2 ± 3.7 years (range from 1 month to 15 years) Table 1.

Table 1: Age distribution of children

Age (Years)	No	(%) Total
Under I year	8	(1.7%)
1-5	233	(49.3%)
6-15	232	(49.0)
Total	473	(100%)

Sex distribution

There were significantly more males (n = 332) than females (n = 141) ($X^2 = 77.1$, P < 0.01). Male: female ratio was 2.4: 1. Mean age of all males was 6.7 ± 3.7 years while that for the females was 5.2 ± 3.5 years. The males were significantly older than the females (P<0.0001).

Etiology of injury

Over 60 percent of all injuries were due to falls and significantly more pre-school children were involved in a fall when compared with the older children 6 years and above ($X^2 = 15.11$, P<0.001). Falls, motor vehicle accidents followed by direct blunt trauma were responsible for over 97 percent of all orofacial injuries in children. Other causes of injury are shown in Table 2.

Table 2: Etiology of injury in ch	1110	irer
-----------------------------------	------	------

Cause	No	% Total
Motor vehicle	90	(60.5%)
Direct blunt trauma	84	(17.8%)
Gun shot	4	(0.8%)
Burns	4	(0.8%)
Unknown	5	(1.1%)
Total	473	100%

For all motor vehicle injuries, none of the 86 children occupants had seat belts on, while the remainders were knocked down as pedestrians.

Site of injury

The commonest site of injury was the forehead region which was followed by the peri-orbital and eye areas and areas around the mouth. Together they accounted for about two-thirds of all injuries (Table 3).

Table 3:	Anatomical	site of	inj	ury
				_

Anatomical site	No	% Total
Forehead	114	(24.1%)
Eyes/Periorbital	110	(23.3%)
Lips/tongues/gums/	105	(22.2%)
dento-alveola/palate		
Chin and Mandible	41	(8.7%)
Nose	39	(8.2%)
Multiple sites	29	(6.1%)
Cheek	26	(5.5%)
Maxilla	9	(1.9%)
Total	473	100%

Seasonality of injuries

Of the 473 cases, 356 (75.3%) took place during the summer months while only 117 (24.7%) occurred during the winter months. The summer months were significantly associated with the occurrence of injuries ($X^2 = 96.1$, P< 0.0001).

Place of injury

The overwhelming majority of injuries in children under the age of 6 years occurred at Home, while the highway/street was the commonest location for injuries in the older children (6 years and above). Figure 1 show the place of injury according to age of the child.



Fig. 1: Location of injury by age

Multiple regression analysis

Logistic regression analysis was used to construct a model of variables to predict the occurrence of injury and to eliminate effect of confounders. Age and seasonality were significant determinants of injury in children and together predicted 46.9 percent in the variance due to injury. Significantly more injuries

Table 4: Adjusted odds of having an injury

Variable	OR*	(95% Confidence interval	p-value
4GE Under 6 years 6 years and above	2.0	(1.3-3.3)	0.003
SEASON Winter summer	0.51 1.0	(0.30 - 0.87)	0.013

*Quasi - Maximum likelihood adjusted odds ratio

occurred in the preschool children (under 6 years) and significantly fewer injuries occurred during the winter months. Sex of the child was not significant following logistic regression analysis (Table 4).

Discussions

Injury and types among children outside developed countries have not received sufficient attention in literature. Thisinformation is needed for health facility planning purposes, to determine treatment plan and for designing appropriate preventive measures to reduce the portion of children involved in such injuries.

From the above data the commonest cause of injury in children was due to a fall. This occurred in nearly two-thirds of thecases and most of the injuries occurred at home. Falls from heights have significant morbidity and mortality risks and most falls have been shown to occur in the home environment The younger children (under 6 years) in particular were more likely to sustain injuries due to falls, this finding compares favorably with that of several authors [2,17]. Various reports blame unnecessary risks during play and insufficient supervision by adults at home for most injuries which occur at home [2,17,19,20]. That information is essentially lacking in the data presented, however, making the home environment safe and directing health education programmes towards the primary caretakers of young children will go a long way towards preventing injuries especially those due to falls in children. Though the highways and road traffic accidents were responsible for most of the injuries in the older children, use of seat belt was generally absent in this study. Use of seat belts is recommended but not mandatory in this community and previous studies have also shown a very low level of use [7]. Few children were knocked down as pedestrians and efforts should also be made to ensure safe driving along the highway.

Seasonality of injury has been noted by some authors, with most injuries occurring in the summer [17]. From the data presented, the summer months were significantly associated with injuries while the winter months were not. During the summer months, the children are at a greater risk of sustaining injuries for two major reasons, one is that schools are usually out on holidays and secondly, the weather is conducive for outdoor recreational activities and sports. Children are more likely to receive close supervision while at school in this community. These factors, in addition to the restrictive winter weather, could be responsible for the fewer injuries recorded in the winter months. This may offer an explanation as to why fewer injuries occurred among the older children, since it is obligatory in this region for children 6 years and above to be in school.

Several earlier studies reported a preponderance of injuries in males [1,10,11,21,23]. This is in agreement with what was found in this study following bivariate analysis. However following logistic regression analysis, the data did not show any significant sex difference in injuries. The large unadjusted effect of the male gender following bivariate analysis may be due to the large number of males seen in this study as well as in the other studies. Several other studies in agreement with this study show no difference in prevalence between boys and girls [12,13]. Following logistic regression analysis, age was a determining factor in this study, and the younger children (under 6 years) were more likely than the older ones (6 years and above) to be injured. This finding is in contrast to what was found in another longitudinal study carried out in Europe [22]. Having a safe home environment, health education programmes directed towards the caretakers of young children and the legislature in support of seat belts especially for minors will go a long way towards

reducing childhood injuries in this community. A multi disciplinary clinical management approach is envisaged due to the different sites of injury and facilities would need to be equipped to take care of all types of oro-facial injuries.

References

- Haug RH, and Foss J. Maxillofacial injuries in the pediatric patient. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2000; 90(2): 126-134.
- Zerfowski M, and Bremerrich A. Facial trauma in chil dren and adolescents. Clin Oral Invest 1998; 2(3): 120-4.
- Grassner R, Bosch R, Tuli T, and Emshoff R. Preva lence of dental trauma in 6000 patients with facial injuries: implications for prevention. Oral Surg Oral Med Oral Pathol Oral Radiol 1999; 87(1): 27-33.
- Harrington MS, Eberthart AB, and Knapp JF. Dentofacial trauma in children. ASDC J Dent Child 1988; 55 (5): 334-338.
- Acton CH, Nixon JW, and Clark RC. Bicycle riding and oral/maxillofacial trauma in young children. Med J Aust 1996; 2: 165(5): 249-251
- Bamjee Y, Lownie JF, Cleaton-Jones PE, and Lownie MA. Maxillofacial injuries in a group of South Africans under 18 years of age. Br J Oral Maxxillofacial Surgery 1996; 34(4): 298-302.
- Lawoyin D.O, Lawoyin J.O, and Lawoyin T.O. Frac tures of the facial skeleton in Tabuk, North West Armed Forced Hospital Saudi Arabia, a five-year review. Afr J Med Med Sci 1996; 25(4): 385-387.
- Mcheik JN, Vergnes P, and Bononny JM. Treatment of facial dog bite injuries in children: a retrospective study. J Ped Surg 2000; 35(4): 580-583.
- Naidoo S. A profile of the oral facial injuries in child physical abuse at a children's hospital. Child Abuse negl. 2000; 24(4): 521-534.
- Borssen E, and Holm AK. Traumatic dental injuries in a cohort of 16 year olds in northern Sweden. Endod Dent Traumatol 1997; 13(6): 276-80.
- Taher AA. Pediatric facial injuries in Tehran: a review of 87 patients. J craniofacial Surg. 1993; 4(1): 21-7.
- Marcenes W, al Beiruff N, Tayfour D., and Issa S. Epidemiology of traumatic injuries to the permanent incisors of 9-12 year olds school children in Damas cus, Syria. Endod Dent Traumatol 1999; 15(3): 117-223
- Otuyemi OD, Segun-Ojo IO, and Adegboye AA. Traumatic anterior dental injuries in Nigerian preschool children. East Afr Med J 1996; 73(9): 604-606.
- Cole B, and Welbury R. Malformation 1 the primary and permanent dentitions following trauma prior to tooth eruption: a case report. Endod Dent Traumatol 1999; 15(6): 294-296.
- Girotto JA, Gamble WB, Robertson B, Redett R, Muchlberser and T, Majer M et al. blindness after reduction of facial fractures. Plast Reconstr Surg 1998; 102 (6): 1821-34 dental disturbances after facial burns and pressure garments. J Burns Care Rehabil 1996; 17(4): 338-345.
- Fricke NB, omnell ML, Dutcher KD, Hollender LG, Engraw CH. Skeletal and dental disturbances after facial burns and pressure garments. J.Burns care Rehabil. 1996; 17 (4): 338 - 345
- Dean A, Dean J, Coulombier D et al. EPI Info version 6. A Word processing, database, statistical program for epidemiology on microcomputer.

Atlanta Georgia. Center for Disease Control and Prevention, 1996.

- SYSTAT 8.0 for Windows. SPSS Inc Chicago, USA 1998.
- Luz JG, and Di Mase F. Incidence of dentoalveolar injuries in hospital emergency room patients. Endod Dent Traumatol 1994; 10(4): 188-190.
- 20. Onetto JE, Flores MT, and Garbarino ML. Dental trauma in children and adolescents in Valparaiso, Chile. Endod Dent Traumatol 1994; 10(5): 223-227.
- Oji C. Fractures of the facial skeleton in children: a survey of patients under the age of 11 years. J cranio maxillofacial surgery; 26(5): 322-325.
- Rohana J, Ong LC, and Abu Hassan A. Epidemiology of head injuries in Malaysian children: a hospital-based study. Med J Malaysia 1998: 53(3): 217-22.
- Denloye OO, Fasola AO, and Arotiba JT. Dental emergencies in children seen at University College Hospital, Ibadan, Nigeria - a five year review. Afr J. Med. Med Sci 1998; 27 (3-4): 197 - 199